



二甲双胍和你的扫描 重要指示

本资料为正在服用含二甲双胍的药物并计划接受造影扫描或造影剂检查的病人给出重要指示。

你的保健医生建议你做 CT 扫描或其他造影检查。作为这种检查的一部分，医生将向你的静脉注入碘造影剂（放射性染料）。


这种造影剂可能会影响你的肾脏将二甲双胍清除出身体的功能。

一些含二甲双胍的常用药物是：

- Fortamet
- Glucophage
- Glucophage XR
- Glumeta
- Riomet

我何时可以开始重新服用我的药物？

- 做完造影扫描后，你需要 **等 48 小时**，然后才能够重新服用你的二甲双胍药物。
- 在重新开始服用你的二甲双胍药物之前，你可能还需要做 **血液检查**。扫描当天，你的保健医生将会给你有关这方面的指示。



如果你有对你的造影检查或你的药物有任何问题，请向你的保健医生咨询。

有任何问题吗？

你的问题非常重要。如果你有任何问题或担忧，请致电你的医生或医疗提供机构。

造影服务部：
206-598-6200

Metformin and Your Scan

Important instructions

This handout gives instructions for patients who take medicine that contains metformin and who are scheduled for an imaging scan or procedure that uses contrast material.

Your health care provider has recommended that you have a CT scan or other imaging procedure. As part of this procedure, *iodine contrast material* (radiology dye) will be injected into your veins.


This contrast material may affect how your kidneys clear metformin from your body.

Some common medicines that contain metformin are:

- Fortamet
- Glucophage
- Glucophage XR
- Glumeta
- Riomet

When can I start taking my medicine again?

- You will need to **wait 48 hours** after you complete your imaging scan before you start taking your metformin medicine again.
- You may also need a **blood test** before you can start taking your metformin medicine again. You will receive instructions about this from your health care provider on the day of your scan.



Please talk with your health care provider if you have any questions about your imaging procedure or your medicines.

QUESTIONS?

Your questions are important. Call your doctor or health care provider if you have questions or concerns.

Imaging Services:
206-598-6200